

2019-01

Compositional response of Amazon forests to climate change.

Esquivel-Muelbert, A

<http://hdl.handle.net/10026.1/12861>

10.1111/gcb.14413

Global Change Biology

Wiley

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Compositional response of Amazon forests to climate change

Running head: Amazon tree community change

Adriane Esquivel-Muelbert¹, Timothy R. Baker¹, Kyle G. Dexter^{2,3}, Simon L. Lewis^{1,4}, Roel Brien¹, Ted R. Feldpausch⁵, Jon Lloyd⁶, Abel Monteagudo-Mendoza^{7,8}, Jos Barlow^{9,10}, Damien Bonal¹¹, Jérôme Chave¹², Nallaret Davila Cardozo¹³, Terry Erwin¹⁴, Sophie Fauset¹, Bruno Hérault^{15, 16}, Susan Laurance¹⁷, Lourens Poorter¹⁸, Lan Qie¹⁹, Clement Stahl²⁰, Martin Sullivan¹, Hans ter Steege^{21,22}, Vincent Antoine Vos^{23, 24}, Pieter A. Zuidema^{18,25}, Everton Almeida²⁶, Edmar Almeida de Oliveira²⁷, Esteban Álvarez Dávila^{28,29,30,31}, Ana Andrade³², Simone Aparecida Vieira³³, Luiz Aragão^{5,34}, Alejandro Araujo-Murakami³⁵, Eric Arets³⁶, Luzmila Arroyo³⁷, Gerardo A. Aymard C.³⁸, Plínio Barbosa Camargo³⁹, Jorcely G. Barroso⁴⁰, Frans Bongers¹⁸, Rene Boot⁴¹, José Luís Camargo³², Wendeson Castro⁴², Victor Chama Moscoso^{8,43}, James Comiskey^{14,44}, Fernando Cornejo Valverde⁴⁵, Antonio Carlos Lola da Costa⁴⁶, Jhon del Aguila Pasquel^{47,48}, Tony Di Fiore⁴⁹, Luisa Fernanda Duque³¹, Fernando Elias²⁷, Julien Engel^{50,51}, Gerardo Flores Llampazo⁵², David Galbraith¹, Emanuel Gloor¹, Rafael Herrera Fernández^{53,54}, Niro Higuchi⁵⁵, Eurídice Honorio Coronado⁵⁶, Wannes Hubau⁵⁷, Eliana Jimenez-Rojas⁵⁸, Adriano José Nogueira Lima⁵⁹, Ricardo Keichi Umetsu²⁷, William Laurance⁶⁰, Gabriela Lopez-Gonzalez¹, Thomas Lovejoy⁶¹, Yadvinder Malhi⁶², Beatriz Marimon²⁷, Ben Hur Marimon Junior²⁷, Omar Aurelio Melo Cruz⁶³, Paulo S. Morandi²⁷, David Neill⁶⁴, Percy Núñez Vargas⁸, Alexander Parada Gutierrez³⁵, Guido Pardo²⁴, Julie Peacock¹, Marielos Peña-Claros^{18,25}, Maria Cristina Peñuela-Mora⁶⁵, Pascal Petronelli⁶⁶, Georgia C. Pickavance¹, Nigel Pitman⁶⁷, Adriana Prieto⁶⁸, Carlos Quesada⁵⁹, Hirma Ramírez-Angulo⁶⁹, Maxime Réjou-Méchain⁷⁰, Zorayda Restrepo Correa⁷¹, Anand Roopsind⁷², Agustín Rudas⁶⁸, Rafael Salomão^{73,74}, Natalino Silva⁷⁵, Javier Silva Espejo⁷⁶, Marcos Silveira⁷⁷, James Singh⁷⁸, Juliana Stropp⁷⁹, John Terborgh⁸⁰, Raquel Thomas⁷², Marisol Toledo³⁵, Armando Torres-Lezama⁸¹, Luis Valenzuela Gamarra⁷, Peter J. van de Meer⁸², Geertje van der Heijden⁸³, Peter van der Hout⁸⁴, Rodolfo Vasquez Martinez⁷, Cesar Vela⁸⁵, Ima Célia Guimarães Vieira⁸⁶, Emilio Vilanova Torre^{69,87}, Oliver L. Phillips¹

¹ School of Geography, University of Leeds, Leeds, LS2 9JT, UK

² Royal Botanic Garden of Edinburgh, EH3 5LR, Edinburgh, UK

³ School of Geosciences, University of Edinburgh, Edinburgh, UK

⁴ Department of Geography, University College London, London, UK

5 Geography, College of Life and Environmental Sciences, University of Exeter, Exeter, UK
 6 Department of Life Sciences, Imperial College London, UK
 7 Jardín Botánico de Missouri
 8 Universidad Nacional de San Antonio Abad del Cusco, Peru
 9 Lancaster Environment Centre, Lancaster University, Lancaster, UK
 10 Museu Paraense Emilio Goeldi, Belém, Brazil
 11 Université de Lorraine, AgroParisTech, INRA, UMR Silva, 54000 Nancy, France
 12 Laboratoire Evolution et Diversité Biologique (EDB), UMR 5174 CNRS/UPS Bâtiment 4R1, 118, route de
 13 Narbonne, 31062 Toulouse, France
 14 Facultad de Ciencias Biológicas, Universidad Nacional de la Amazonía Peruana, Peru
 15 Smithsonian Institution, 1100 Jefferson Drive, Washington DC 20560
 16 Cirad, Univ Montpellier, UR Forests & Societies, Montpellier, France.
 17 INPHB, Institut National Polytechnique Félix Houphouët-Boigny, Yamoussoukro, Ivory Coast.
 18 Centre for Tropical Environmental and Sustainability Science (TESS) and College of Marine and Environmental
 19 Sciences, James Cook University
 20 Wageningen University and Research, Forest Ecology and Forest Management group, Wageningen, The Netherlands
 21 Department of Life Sciences, Imperial College London, Silwood Park Campus, Ascot, SL5 7PY, UK
 22 INRA, UMR EcoFoG, AgroParisTech, CNRS, Cirad, Université des Antilles, Université de Guyane, 97310 Kourou,
 23 France
 24 Naturalis Biodiversity Center, PO Box 9517, Leiden, 2300 RA, The Netherlands
 25 Systems Ecology, Free University, De Boelelaan 1087, Amsterdam, 1081 HV, Netherlands
 26 Centro de Investigación y Promoción del Campesinado - Norte Amazónico, Riberalta, Bolivia
 27 Universidad Autónoma del Beni, Riberalta, Bolivia
 28 Programa Manejo de Bosques de la Amazonía Boliviana
 29 Instituto de Biodiversidade e Floresta, Universidade Federal do Oeste do Pará, Pará, Brazil
 30 Universidade do Estado de Mato Grosso, Campus de Nova Xavantina, Brazil
 31 Interconexión Eléctrica S.A
 32 Red para la Mitigación y Adaptación al Cambio Climático de la UNAD- Red MiA
 33 Servicios Ecosistémicos y Cambio Climático, Jardín Botánico de Medellín
 34 Socioecosistemas y clima sostenible, Fundación con Vida
 35 Instituto Nacional de Pesquisas da Amazônia, Projeto Dinâmica Biológica de Fragmentos Florestais, Manaus, Brazil
 36 Universidade Estadual de Campinas, Campinas, São Paulo, Brazil
 37 National Institute for Space Research (INPE), São José dos Campos-SP, Brazil
 38 Museo de Historia Natural Noel Kempff Mercado, Universidad Autónoma Gabriel René Moreno, Santa Cruz, Bolivia
 39 PROMAB, Riberalta, Bolivia
 40 Universidad Autónoma Gabriel Rene Moreno, El Vallecito, Santa Cruz-Bolivia
 41 UNELLEZ-Guanare, Programa de Ciencias del Agro y el Mar, Herbario Universitario (PORT)
 42 Universidade de São Paulo, São Paulo, Brazil
 43 Universidade Federal do Acre, Campus de Cruzeiro do Sul, Acre, Brazil
 44 31 Tropenbos International and Group Ecology and Biodiversity, Beta Faculty, Utrecht University, The Netherlands
 45 Programa de Pós-Graduação Ecologia e Manejo de Recursos Naturais, Universidade Federal do Acre, Acre, Brazil
 46 Jardim Botânico de Missouri
 47 Inventory and Monitoring Program, National Park Service, 120 Chatham Lane, Fredericksburg, VA 22405, USA
 48 Andes to Amazon Biodiversity Program, Peru
 49 Universidade Federal do Pará, Pará, Brazil
 50 Instituto de Investigaciones de la Amazonia Peruana
 51 Michigan Tech University
 52 University of Texas at Austin
 53 AMAP, IRD, CIRAD, CNRS, Université de Montpellier, INRA, Boulevard de la Lironde, TA A-51/PS2, F-34398
 54 Montpellier Cedex 5, France
 International Center for Tropical Botany, Department of Biological Sciences, Florida International University, 11200
 SW 8th Street, Miami, FL 33199, USA
 Universidad Nacional Jorge Basadre de Grohmann (UNJBG), Peru
 Centro de Ecología, Instituto Venezolano de Investigaciones Científicas
 ReforeST Group, DIHMA, Universidad Politécnica de Valencia

- 55 Instituto Nacional de Pesquisas da Amazônia - Coordenação de Pesquisas em Silvicultura Tropical, Manaus, Brazil
- 56 Instituto de Investigaciones de la Amazonía Peruana
- 57 Royal Museum for Central Africa, Leuvensesteenweg 13, 3080 Tervuren, Belgium
- 58 Grupo de Investigación en Temas Agroambientales INTEGRA, Tecnológico de Antioquia Institución Universitaria
- 59 Instituto Nacional de Pesquisas da Amazônia, Manaus, Brazil
- 60 Centre for Tropical Environmental and Sustainability Science (TESS) and College of Science and Engineering, James Cook University
- 61 Center for Biodiversity and Sustainability at George Mason University, Fairfax, VA 22030, USA
- 62 Environmental Change Institute, School of Geography and the Environment, University of Oxford
- 63 Universidad de Tolima, Colombia
- 64 Universidad Estatal Amazónica, Puyo, Pastaza, Ecuador
- 65 Universidad Regional Amazónica ikiam, Tena, Ecuador
- 66 Cirad UMR Ecofog (AgrosParisTech, CNRS, INRA, Univ Guyane); Campus agronomique; 97310 Kourou, French Guiana
- 67 Science and Education, The Field Museum
- 68 Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Colombia
- 69 Universidad de los Andes, Venezuela
- 70 Université Paul Sabatier CNRS, UMR 5174 Evolution et Diversité Biologique
- 71 Servicios Ecosistémicos y Cambio Climático (SECC), Fundación Con Vida & Corporación COL-TREE
- 72 Iwokrama International Centre for Rainforest Conservation and Development
- 73 Pós-Graduação em Botânica Tropical, Universidade Federal Rural da Amazônia/Museu Paraense Emílio Goeldi, Pará, Brazil
- 74 Programa de Pós-Graduação em Agricultura e Ambiente, Universidade Estadual do Maranhão, São Luís, Maranhão, Brazil
- 75 Serviço Florestal Brasileiro
- 76 Departamento de Biología. Universidad de La Serena. Casilla 554 - La Serena, Chile
- 77 Universidade Federal do Acre, Museu Universitário, Acre, Brazil
- 78 Guyana Forestry Commission, Georgetown, Guyana
- 79 Joint Research Centre of the European Commission
- 80 Center for Tropical Conservation, Nicholas School of the Environment, Duke University
- 81 Universidad de Los Andes, INDEFOR, Mérida, Venezuela
- 82 Van Hall Larenstein University of Applied Sciences, Forest and Nature Management Group
- 83 University of Nottingham, Nottingham, UK
- 84 van der Hout Forestry Consulting, Rotterdam, the Netherlands
- 85 Facultad de Ciencias Forestales y Medio Ambiente, Universidad Nacional San Antonio Abad Cusco, Puerto Maldonado, Peru
- 86 Museu Paraense Emílio Goeldi, Pará, Brazil
- 87 School of Environmental and Forest Sciences, University of Washington, Seattle, USA.

29

30 **Corresponding author:** Adriane Esquivel-Muelbert

31 Telephone: +44 (0) 113 34 33907

32 Email: adriane.esquivel@gmail.com

33 **Key words:** bioclimatic niches; climate change; tropical forests; functional traits; temporal
34 trends; compositional shifts.

35 **Paper type:** Primary Research

Abstract

Most of the planet's diversity is concentrated in the tropics, which includes many regions undergoing rapid climate change. Yet, while climate-induced biodiversity changes are widely documented elsewhere, few studies have addressed this issue for lowland tropical ecosystems. Here we investigate whether the floristic and functional composition of intact lowland Amazonian forests have been changing by evaluating records from 106 long-term inventory plots spanning 30 years. We analyse three traits that have been hypothesized to respond to different environmental drivers (increase in moisture stress and atmospheric CO₂ concentrations): maximum tree size, biogeographic water-deficit affiliation and wood density. Tree communities have become increasingly dominated by large-statured taxa, but to date there has been no detectable change in mean wood density or water deficit affiliation at the community level, despite most forest plots having experienced an intensification of the dry season. However, among newly recruited trees dry-affiliated genera have become more abundant, while the mortality of wet-affiliated genera has increased in those plots where the dry season has intensified most. Thus, while changes in compositional dynamics (recruits and mortality) are consistent with climate-change drivers, these have yet to significantly impact whole-community composition. Overall, we find that the increase in atmospheric CO₂ appears to be driving a shift within tree communities to large-statured species and that climate changes to date will eventually impact Amazon forest composition, but long generation times of tropical trees mean that the speed of the floristic response is lagging behind the rate of climatic change.

Introduction

Tropical forests represent the world's most biodiverse ecosystems, as well as providing its largest stores of living carbon and contributing more to biomass productivity than any other biome on the planet. Changes here can therefore have global consequences, potentially nowhere more so than in Amazonia where between 6,000 and 16,000 tree species exist (Cardoso *et al.*, 2017, ter Steege *et al.*, 2013) and as much as 100 Pg of carbon is stored in biomass (Feldpausch *et al.*, 2012). While the physical, chemical, and biological environment have all been changing over recent decades, it is the changes in climate – both documented and projected – which are widely expected to cause some of the most profound changes in forest communities and ecosystem processes (Esquivel-Muelbert *et al.*, 2017a, Thomas *et al.*, 2004). For example, higher temperatures and intensifying drought may threaten larger trees due to hydraulic failure (McDowell & Allen, 2015, Rowland *et al.*, 2015), which could eventually compromise forest biomass and productivity. In Amazonia, because forest diversity is concentrated in the wetter, least seasonal forests (Francis & Currie, 2003, Gentry, 1988), a persistent lengthening of the dry season might threaten a large portion of tropical biodiversity. Here we aim to better understand how Amazonian trees have responded to the last 30 years of environmental change, by analysing floristic records from long-term tree monitoring in the Neotropics to assess the potential compositional changes to date.

Changes in biodiversity attributed to climate change have already been documented in a wide variety of ecosystems (e.g. Bowler *et al.*, 2017, Chen *et al.*, 2009), including in some tropical locations, but so far there is remarkably little evidence of any widespread impacts of climate change on the composition of tropical ecosystems which harbour much of the planet's diversity (Duque *et al.*, 2015, Fauset *et al.*, 2012, van der Sande *et al.*, 2016). In contrast, there

is evidence for large-scale changes in the structure (i.e., aboveground biomass) and dynamics (e.g., mortality and productivity) of old-growth tropical forests. In many forests, apparently undisturbed by humans, both aboveground biomass and the rate of ecological processes such as growth and recruitment have increased (Brienen *et al.*, 2015, Lewis *et al.*, 2009b, Phillips & Gentry, 1994, Qie *et al.*, 2017), while in Amazonia increases in mortality have caused a recent weakening of the biomass carbon sink (Brienen *et al.*, 2015). However, it remains unclear whether these structural and dynamic changes are also associated with concerted changes in the species richness and composition of Amazonian forests.

In Amazonia, as elsewhere, climate change is a potential leading driver of changes to the ecosystem. During the last few decades, the climate of Amazonia has become more extreme – the length of the dry season and its intensity have increased, whilst precipitation has become more intense during the wet season (Gloor *et al.*, 2015, Hilker *et al.*, 2014). Extreme climate events in recent years include the three strong droughts within a decade (Erfanian *et al.*, 2017, Jiménez-Muñoz *et al.*, 2016, Lewis *et al.*, 2011, Marengo *et al.*, 2008, Marengo *et al.*, 2011) and several large-scale episodes of extreme rainfall (Espinoza *et al.*, 2014, Marengo & Espinoza, 2016). In addition to the repeated drought events, precipitation has declined in the south and south east of the basin (25% reduction in rainfall between 2000 and 2012) (Hilker *et al.*, 2014) and higher temperatures are likely to have intensified seasonal evaporative stress across the basin (Jiménez-Muñoz *et al.*, 2013). These changes are consistent with model-based predictions (Duffy *et al.*, 2015), implying that the Amazon may already have entered a new regime of a hotter, more variable climate. The forest has clearly responded to these recent fluctuations in climate – for example, tree mortality rates increased markedly during and after drought events causing at least temporary losses of standing biomass (Brienen *et al.*, 2015, Feldpausch *et al.*, 2016, Phillips *et al.*, 2009, Zuleta *et al.*, 2017). The 2010 drought also

impacted on the basin-wide exchange of carbon dioxide between the vegetation and the atmosphere, with the vegetation becoming a net source of CO₂ during 2010 (Gatti *et al.*, 2014).

In addition to the changes in climate, atmospheric CO₂ concentrations have increased globally from 340 ppm to over 400 ppm over the past 50 years (Conway & Tans, 2016). Carbon dioxide is also a fundamental resource for photosynthesis and higher concentrations are expected to stimulate plant growth (Lloyd & Farquhar, 1996). Indeed, the increase in atmospheric concentrations of CO₂ is a potential driver of the observed increase in aboveground biomass and turnover rates in tropical forests (Brienen *et al.*, 2015, Lewis *et al.*, 2009a, Pan *et al.*, 2011, Zhu *et al.*, 2016). Additionally, under higher CO₂ concentrations plants may increase their water-use efficiency with less water being required per unit of carbon fixed. Thus, by allowing plants to use less water, higher CO₂ concentrations could alleviate the effect of increasing aridity (Lloyd & Farquhar, 2008, van der Sleen *et al.*, 2015).

Interpreting potential shifts in tropical floristic composition and linking them with possible drivers is a considerable challenge due to the very high diversity of tropical forests and their large spatial extent. However, if sufficient high-quality, long-term floristic monitoring is available, then the approach of analysing shifts in a suite of functional traits to describe how communities change over time can be used to link floristic changes to their drivers (McGill *et al.*, 2006, Violle *et al.*, 2014). For tropical tree species, two largely independent trait axes may have value in addressing these questions. One axis, the life-history trade-off between growth and mortality, can be represented by wood density as it is negatively correlated to growth and mortality rates. This is a reflection of slow-growing trees tending to invest more in wood structure (Chave *et al.*, 2009, Muller-Landau, 2004, Nascimento *et al.*, 2005, Poorter *et al.*,

2010). The other axis is related to the potential size that taxa can attain, representing the capacity of taxa to compete for light (Falster & Westoby, 2005, Poorter *et al.*, 2005).

Environmental changes could have different impacts along each of these ecological axes (Chave *et al.*, 2008, Lewis *et al.*, 2009a). Most notably, with more extended or intense periods of soil water deficit, large trees and those with low wood density may be at greatest risk of hydraulic failure due to cavitation (McDowell & Allen, 2015, Rowland *et al.*, 2015). Large trees have been shown to be particularly affected by artificially-imposed drought (McDowell & Allen, 2015, Nepstad *et al.*, 2007, Rowland *et al.*, 2015) and drought events (Bennett *et al.*, 2015, Phillips *et al.*, 2010). On the other hand, several observations from tropical forests show a decline of small understory taxa associated with increases in soil water deficit increases (e.g. Condit *et al.*, 1996, Enquist & Enquist, 2011, Fauset *et al.*, 2012, Feeley *et al.*, 2011a). To explain these observations, it has been hypothesized that small taxa have shallower roots and are therefore be more vulnerable compared to large, deep rooted trees to long-term drying trends (Condit *et al.*, 1996, Fauset *et al.*, 2012). Although the link between rooting depth and tree size is still unclear (Stahl *et al.*, 2013), this hypothesis is consistent with wetter areas tending to have more densely populated understoreys (Malhi *et al.*, 2002, Pitman *et al.*, 2002) and taller forests being less sensitive to precipitation variability (Giardina *et al.*, 2018). The on-going increase in atmospheric carbon dioxide is also expected to cause changes in species composition as it is predicted to favour those trees that have greater competitive capacity to access light (Coomes *et al.*, 2011, Laurance, 2004), consequently increasing the mean potential tree size within the community (e.g. Laurance *et al.*, 2004) and to favour fast-growing trees, potentially leading to communities with lower wood density.

Given the uncertainty about how tree size relates to responses to a drying climate, the geographic distributions of species over precipitation gradients offer an alternative source of information with which to predict and to infer the effects of drought on floristic composition. The spatial distribution of tree taxa over precipitation gradients has been shown to provide a valuable proxy for drought-tolerance in observational studies and experiments (Engelbrecht *et al.*, 2007, Esquivel-Muelbert *et al.*, 2017b). If drought is increasingly affecting Amazonian forests, we might therefore expect concerted shifts in tree communities towards more dry-affiliated components. A compositional shift towards more dry-tolerant taxa as a consequence of an increase in moisture-stress has been reported locally for sites in southern Ghana (Fauset *et al.*, 2012), Central America (Enquist & Enquist, 2011, Feeley *et al.*, 2011a), and parts of the Amazon (Butt *et al.*, 2014).

Here, we aim to quantify the shifts in the floristic composition of Amazonian tree communities, and test the hypothesis that recent climatic drying is already impacting the composition of Amazonian forests. The dataset derives from 106 long-term, tree inventory plots distributed across intact closed-canopy moist forests in lowland Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru, and Venezuela (Appendix S1). We analyse monitoring records from as early as 1985 onwards to as recently as 2015, deliberately excluding any possible influence of the 2015-16 El Niño drought from our analysis. We investigate trends within the overall community composition, as well as among recruits, trees that died, and in the growth rate that occurred within each census interval (Figure 1). We analyse compositional shifts along these three trait axes, which we demonstrate to be independent: life-history (using wood density as a proxy), potential size, and bioclimatic distribution (Appendix S2). Based on predictions from plant physiology supported by experimental studies, we expect increases in dry season duration or intensity to shift floristic composition towards dry-affiliated and taxa

with smaller-statured genera with high wood density (McDowell & Allen, 2015, Rowland *et al.*, 2015). Additionally, we examine trends in abundance for individual genera, which allows us to understand which taxa dominate the changes in functional composition.

Methods

Field observations and forest dynamics

We investigate the trends in functional and floristic composition of tree communities by analysing long-term data from permanent tree inventory plots in the Amazon and adjacent lowland forests (Appendix S1). A total of 106 South American forest plots from the RAINFOR network (Malhi *et al.*, 2002), accessed via the ForestPlots.net repository (Lopez-Gonzalez *et al.*, 2011), met our criteria: (i) sampling lowland (< 1000 m.a.s.l), *terra firme*, intact, moist forests (i.e., where mean maximum cumulative water deficit (MCWD) over the last century is less negative than -300 mm y⁻¹); (ii) having been monitored throughout the period of the two intense dry seasons in 2005 and 2010 (see Supplementary material for analyses including plots that were not monitored during 2005 and 2010); (iii) having had regular monitoring, thus excluding any plots where census interval lengths differed by more than 10 years; (iv) having at least 80% of tree stems identified at least to genus level. The selected plots have a mean area of 1.25 ha (95% CI = 1.16, 1.35), and have been re-censused on average 7 times, with a mean census interval of 2.8 years.

The long-term plots were monitored following a standardized protocol (Phillips *et al.*, 2016). Full methodological details are given elsewhere (Brienen *et al.*, 2015). In brief, all trees and palms ≥ 10 cm diameter (D) at 1.3 m (or above-buttress) are tagged, identified to the species level (when possible), have their D measured, and the point of measurement marked and

recorded. At every census, trees previously recorded are re-measured, and new recruits - i.e., trees that have newly attained 10 cm when the plot is revisited - tagged, measured and identified, and notes are taken about the individuals that died between censuses. Lianas and non-woody freestanding plants (*Phenakospermum*) are excluded from our analyses.

Climate data

To characterise the change in moisture-stress we calculated temporal trends in maximum cumulative water deficit (MCWD - Aragao *et al.*, 2007) for each plot. MCWD represents the most negative value of water deficit (wd), i.e., the difference between precipitation (P) and evapotranspiration (E) within each year, where for each month (n) wd is quantified as:

$$\begin{aligned} & \text{If } wd_{n-1} - E_n + P_n < 0; \\ & \text{then } wd_n = wd_{n-1} - E_n + P_n \quad \text{Eq (1)} \\ & \text{else } wd_n = 0 \end{aligned}$$

In other words, MCWD is an annual water deficit metric which takes into account both the length and intensity of the dry season based solely on climatic variables, i.e., ignoring soil properties. The calculation of MCWD does not necessarily follow the calendar year, as for tropical forests in the northern hemisphere the annual dry season typically spans two calendar years (between October and March). Thus, the starting point, i.e., when $n = 1$, was defined climatologically as the wettest month in the first year in the time series (i.e., 1985), rather than the first month of that calendar year.

In addition to annual MCWD, following Feldpausch *et al.* (2016), we also estimated the intensity of the most extreme dry season between two consecutive censuses, hereafter termed MCWD_i. This metric represents a measure of maximum environmental disruption between two censuses, i.e., the most negative value of annual MCWD between each successive pair of censuses. Only complete years were considered for this calculation. For the first census of each plot the MCWD_i was calculated as the most negative MCWD values within the 3-year period preceding that census. This time window is equivalent to the mean census interval within the data (2.8 years).

Climate data were obtained from the Climatic Research Unit (CRU), at 0.5° spatial resolution from 1985 to 2015 (Harris *et al.*, 2014) where evapotranspiration is calculated based on the Penman–Monteith equation (Allen *et al.*, 1994) using information on temperature, cloud cover and vapour pressure (Harris *et al.*, 2014). MCWD trends calculated from this ground-based precipitation data source were consistent with satellite-based data from the Tropical Rainfall Measuring Mission (TRMM - Huffman *et al.*, 2007) at 0.25° spatial resolution between 1998 and 2010 ($R^2 = 0.87$; $P < 0.0001$ - see Appendix S3 for more detail). Here we opt to use ground-based data from CRU as this covers the whole time window of the floristic analyses.

Traits

We describe Amazonian tree genera in terms of the three basic traits shown in previous work and in Appendix S2 for our data to represent largely independent axes of fundamental plant characteristics and each potentially responsive to environmental change: **Potential Size (PS) (cm)**: the 95th percentile of the distribution of trees' diameter derived from a set of 530 inventory plots across the Amazon Basin (Fauset *et al.*, 2015); **Water deficit affiliation (WDA) (mm)**: derived from relative abundances across 513 inventory plots distributed along

a large gradient of MCWD across the Western Neotropics (Esquivel-Muelbert *et al.*, 2017a). This metric quantifies the preferred dry season severity for each taxon and is calculated as the dry season severity (measured as MCWD) where a taxon occurs weighted by the taxon's relative abundance in each location (Esquivel-Muelbert *et al.*, 2017a). Taxa affiliated to dry conditions show the most negative values of WDA, whilst the most wet-affiliated ones have values of WDA equal to zero; **Wood density (WD) (g cm^{-3})**: values were extracted from the Wood Density Database (Chave *et al.*, 2009, Zanne *et al.*, 2009). There was no indication of any pairwise relationship among any of the traits: WDA vs. PS ($R^2 = 8 \times 10^{-3}$) WDA vs. WD ($R^2 = 5 \times 10^{-4}$) and WD vs. PS ($R^2 = 2 \times 10^{-3}$).

The diversity of Amazonian flora hinders us from performing consistent species-level analyses as comprehensive trait data are still missing for the large majority of Amazonian tree species. Therefore, our analyses were performed at the genus-level and the mean trait value of the genus was assigned to each individual. Our approach is expected to be adequate to capture the actual functional shifts in Amazonia as it captures most of the trait variation of Amazonian trees, which is mostly accounted for at the family and genus level (see Baker *et al.*, 2004, Coelho de Souza *et al.*, 2016, Fyllas *et al.*, 2009, Patiño *et al.*, 2009). Although genus-level analyses still of course miss some information on trait variation, analyses at this taxonomic resolution are potentially more conservative with respect to the hypothesis of environmental-driven floristic change than analyses at the species level as they use mean genus-level values instead of (more extreme) species-level values. Results of analyses using species-level traits, for those taxa where data are available, do not differ from genus-level results (see Appendix S8).

Genus-level trait data were missing for 6%, 9% and 0.04% of all stems for PS, WDA and WD respectively. For these cases, the mean trait values from the family were used, following established conventions (Baker *et al.*, 2004, Flores & Coomes, 2011) and considering the phylogenetic conservatism of PS and WD for Amazonian trees (Coelho de Souza *et al.*, 2016). For the small proportion of individuals belonging to families for which no trait information was available, we used average trait values of the community in the same census was used. For those stems not identified to family level (3.9%), we applied the mean trait for all dicot individuals of the plot-census during which the tree was recorded. Considering the low proportion of missing data the imputation technique is unlikely to strongly affect our results (Penone *et al.*, 2014).

Then, to obtain a census-level value for each of the three traits, we scaled the genus-level traits to the community level by calculating the community-weighted mean (CWM sensu Diaz *et al.*, 2007) for each trait in each census. For each of the 743 censuses across 106 plots, the CWM of each of these components was calculated as the mean trait value across the genera of the community, weighted by (1) the number of stems and (2) the total basal area occupied by each genus. Community-weighted means were calculated for the whole community and for each component of forest dynamics, i.e., the recruits (new trees that reach the 10 cm D cut-off after each census interval), losses (those trees that died within each census interval) and the basal area gain of those trees that survived the census interval (Figure 1).




Figure 1 – Schematic model representing the different components of forest demography. The box on the left represents an inventory plot of a forest community at the first census (C_{t0}), whilst the box on the right shows the community at the second census (C_{t1}). At C_{t1} recruits (R), i.e., those trees that attained 10 cm of diameter within the census interval, will now be part of the community analysed. Other trees will have died thus leaving the community, here called *losses* (L). Those trees from C_{t0} that survive through the census interval are expected to grow (G). Thus, the basal area of the whole community at t_1 is $C_{t1} = C_{t0} + G_{t1} + R_{t1} - L_{t1}$ and the net flux between t_0 and t_1 is $G_{t1} + R_{t1} - L_{t1}$. Here we investigate the trends in the characteristics and identity of genera within these components of forest demography. This figure represents dynamics in basal area terms; similar logic can be applied for stem-based analyses. Note that in this case we would not be interested in the growth of trees surviving from t_0 to t_1 , and so the net flux would be represented as $R_{t1} - L_{t1}$.

Analytical approach

We investigated changes in functional (mean potential size, water deficit affiliation and wood density) and floristic composition (relative abundance of individuals within different genera) by assessing trends over time of these quantities for each plot and their Amazon-level mean.

Trends in functional composition

We investigated functional trends for the different components of forest demography (i.e., whole tree community, recruits and trees that died - Figure 1). These functional trends

were quantified using (1) the bootstrapped mean and 95% CI of plot level linear regression slopes of the community weighted mean (CWM) as a function of time, averaged across all plots, and (2) linear mixed effect models (LMM) of CWM as a function of time with plot identified as a random effect, using function *lmer* from the R package *lme4* (Bates *et al.*, 2015). Whilst the first approach provides information of the overall mean trend across all plots, in which the uncertainty estimate is derived using a non-parametric approach, the second approach gives an estimate of the trend over the whole period (1985-2015), accounting for potential changes in which plots are analysed over different time windows along the 30-year period.

The bootstrapped mean and confidence intervals were calculated from the linear slopes of CWM (i.e., mean plot level traits) for each plot where the CWM in each census (*j*) is a function of the date when the census took place (time):

$$CWM_j \sim \beta_1 + \beta_2 * time_j \quad (Eq\ 2)$$

Then, an Amazon-wide weighted mean and the 95% CI were estimated by randomly resampling values of the plot-level slopes (β_2 from Eq. 2) across all plots, 10,000 times. We further estimate the potential influence of spatial autocorrelation on our results by testing the correlation between the Euclidean distance in the trends in CWM and the spatial distance between the plots (Appendix S7).

The variation in plot area and monitoring period may be expected to influence plot-level trends as smaller plots and those monitored for shorter periods are more likely to be affected by stochastic phenomena, such as tree falls (Fisher *et al.*, 2008). An empirical investigation of the impact of sampling effort (i.e., length of monitoring period and plot size)

on our estimate showed that the variance is expected to scale as a function of the square root of plot area multiplied by the monitoring period (Appendix S4). Thus, to account for the noise attributed to sampling effort we used the square root of plot area multiplied by the monitoring period as weights when calculating Amazon-level mean and confidence intervals of the CWM slopes.

The LMM follows the same logic of the previous analyses with CWM as a linear function of time:

$$CWM_{ij} \sim (\beta_1 + a_i) + (\beta_2 + b_i) * time_{ij} + \varepsilon_{ij} \quad (\text{Eq 3})$$

where CWM in plot i and census j is a function of the date when the census occurred (time). Here time is used as a fixed effect and the random components of the model are the intercept (a) and slope (b) for each plot and the overall residuals (ε). The slope and intercept of each plot were included as random effects considering that both the initial trait value (represented by the plot intercept, a_i) and the intensity of change (represented by the plot slope, b_i) may differ across plots. An exploration of the influence of sampling effort on the variance of the residuals shows no weighting procedure to be required for this analytical framework (Appendix S4).

The influence of climate on functional composition

To explore the potential impact of climate changes on functional change, we tested whether changes in the community are related to changes in climate. First, we calculated the linear trend in MCWD and MCWD_i over the inventory period for each plot. We then used Kendall's τ coefficient to test for the correlation between linear slopes of change in CWMs

(composition) and MCWD (climate) for each plot. For the cases where the correlation was significant, we fit an OLS linear regression with the trends in CWM as a function of trends in MCWD.

Trends in floristic composition

We investigated the influence of individual taxonomic groups on any functional shifts by analysing the relationship between shifts in abundance and trait values. Trends in relative abundance for each taxon were calculated following the same procedure applied above to analyse CWM changes, i.e., by calculating the bootstrapped mean of linear trends and applying LMM, but here using the relative abundance of each genus as a response variable instead of CWM. The bootstrap means were calculated from a modified version of Eq 2 applied to every plot:

$$RA_j \sim \beta_1 + \beta_2 * time_j \quad (Eq\ 4)$$

where RA is the relative abundance of a genus at plot i and census j . We then use the LMM approach by modifying Eq. 3 to:

$$RA_{ij} \sim (\beta_1 + a_i) + (\beta_2 + b_i) * time_{ij} + \varepsilon_{ij} \quad (Eq\ 5)$$

An Amazon-wide slope using each of the methods was calculated for each taxon. The LMM analysis was restricted to genera that occurred in three or more plots. As for the functional analyses, each plot in the bootstrap mean following Eq 4 was weighted by the square root of plot area multiplied by the monitoring period and no weighting procedure was applied for the LMM approach (Eq 5). Trends in abundance were also calculated for families and species (Appendix S12).

Next, we investigated which genera contribute most to the significant functional changes that were detected. When trends in functional composition were significantly different from zero (see *Trends in functional composition* for details), we estimated Kendall's τ coefficient of correlation between Amazon-wide slopes (calculated for the whole community, for recruits, and for losses) for each genus and their trait values (WDA, WD or PS). To ensure that abundance trends were estimated with reasonable levels of uncertainty, we restricted the investigation of the trends in abundance vs. traits relationship to only include genera within the upper 20% quantile of abundance across the whole dataset.

Finally, we calculated the trends in abundance for four Amazonian functional types defined by Fyllas *et al.* (2012) using trait information from a subset of the Amazonian species included in this study. The four Amazonian functional groups (small pioneers, small-statured non-pioneers, tall pioneers and tall non-pioneers) are based on variety of foliar and structural traits (Appendix S10) independent from the traits considered in our main analyses. This therefore provides complementary information to our main analyses.

Results

Climate trends

The forests analysed here have on average experienced a strengthening of the dry season, as measured by maximum cumulative water deficit (MCWD). The plot-level annual MCWD became more negative by on average 1.1 mm y^{-1} (95% CI = 1.4, 0.9) since 1985 (Fig. 2). This represents a marked increase in MCWD across our plots by on average 36% between 1985 and 2014. There are exceptions to the overall drying trend, with some plots near the Andes becoming wetter during this period (Fig. 2).




Figure 2 – Trends in maximum cumulative water deficit (MCWD) across the Amazon Basin. (a) Frequency of annual linear trends in MCWD per plot between 1985 and 2014. Note that for most plots the climate has significantly shifted towards more negative MCWD values. Mean change and 95% confidence intervals (black solid and dashed lines) across our plots weighting plots by plot area were calculated using a bootstrap procedure by resampling the trends in MCWD from all plots 10,000 times with replacement. **(b)** Distribution of annual linear trends in MCWD per plot. Arrows pointing down (in red) represent locations where MCWD has become more negative over time, i.e., the dry season has become more intense. Arrows pointing up (blue) represent less negative values of MCWD meaning that moisture stress decreased. The intensity of the colors in (a) and (b) represent the strength of the climate trend. Note the difference in scale between drying and wetting trends color bars. **(c)** Mean annual MCWD across plots, and 95% CI from resampling among all plots, note lower MCWD values at 2004-2005 and 2009-2010 (grey-shaded rectangles).

Trends in functional composition

Overall, there has been a significant increase in the potential size (PS) of tree communities: the community weighted mean (CWM) of PS when weighted by basal area increased by 0.03 cm y^{-1} (95% CI = 0.02, 0.05 cm y^{-1}) or $0.06\% \text{ y}^{-1}$ with basal gains increasing in PS by 0.06 cm y^{-1} (95% CI = 0.01, 0.12 cm y^{-1}) meaning that plots have been progressively occupied by larger statured genera (Table 2). The increase in basal area-weighted community-level PS holds regardless of the analytical technique, i.e., bootstrapped means or LMM (Table 2, Appendix S6).

For the recruits, dry-affiliated genera have become more abundant, with water-deficit affiliation (WDA) becoming significantly more negative for this segment of the community by 0.45 mm y^{-1} (CI 95% -0.9, -0.03 mm y^{-1}) or $0.3\% \text{ y}^{-1}$ (Table 1). This trend was marginally significant across the basin when calculated using LMM (Appendix S6.1), however it was observed for 62 out of 102 plots ($P=0.03$, two-tailed binomial test) and persisted when our analyses were repeated across a larger dataset (Appendix S6.2) and when grouping together nearby plots to account for spatial autocorrelation in the distribution of plots (Appendix S7.2).

Among the trees that died within the study period (losses), basal area-weighted wood density (WD) decreased by $1 \times 10^{-3} \text{ g cm}^3 \text{ y}^{-1}$ or $0.16\% \text{ y}^{-1}$ in the LMM analyses (Appendix S6). This trend, however, was only observed for 53 out of 102 communities, meaning that positive and negative slopes are equally likely to be observed ($P=0.8$, two tail binomial test), thus it was only marginally significant when using bootstrapped means: $-1 \times 10^{-3} \text{ g cm}^3 \text{ y}^{-1}$ (90% CI = -2×10^{-3} , $-5 \times 10^{-6} \text{ g cm}^3 \text{ y}^{-1}$). No equivalent significant abundance-based trend for losses in WD terms was observed (Table 1, Appendix S6).




Figure 3 – Trends in functional composition between 1985 and 2015 across Amazonia. The y-axes show stem-based community weighted mean (CWM) trends in (a-c) water deficit affiliation (WDA), (d-f) potential size (PS) and (g-i) wood density (WD) at genus level. Values are standardized with respect to the whole plot population to allow comparison among traits meaning that the value for the whole community in the first census is equal to 1. CWM trends are shown for the whole community (a, d, g), recruits (b, e, h), and losses (c, f, i). Grey line and grey-shaded area show standardized mean and 95% CI census-level CWM, which can be influenced by some switching of plots assessed through time. Trend lines show linear mixed models (LMM) considering the slope and intercept of plots as random effects. Slope values for LMM are shown in each graph - these are not standardized by plot population and are shown at a different scale for each trait.

Table 1 – Mean linear slopes in stem-based functional composition in Amazonia. For each trait, we show the bootstrap mean annual changes in community weighted mean (CWM) and 95% confidence intervals (CI, in brackets) weighted by the product of the squared root of plot size and monitoring period. CWM is calculated using: water deficit affiliation (WDA), potential size (PS) and wood density (WD). The analyses were repeated for recruits, losses and the difference between recruits and dead trees (net fluxes). Significant trends are in bold, i.e., where 95% CIs do not overlap zero.

Community	Potential size (cm y ⁻¹)	Water Deficit affiliation (mm y ⁻¹)	Wood Density (g cm ⁻³ y ⁻¹)
Whole community	0.01 (-0.002 0.01)	0.01 (-0.03 0.04)	-1x10 ⁻⁵ (-9x10 ⁻⁵ 6x10 ⁻⁵)
Gains (recruits)	0.07 (-0.03 0.2)	-0.45 (-0.9 -0.03)	-3x10 ⁻⁴ (-2x10 ⁻³ 1x10 ⁻³)
Losses	0.1 (-0.01 0.2)	-0.1 (-0.6 0.3)	2x10 ⁻⁴ (-7x10 ⁻⁴ 1x10 ⁻³)
Net fluxes	-0.03 (-0.2 0.1)	-0.45 (-1 0.1)	-7x10 ⁻⁴ (-2x10 ⁻³ 8x10 ⁻⁴)

Table 2 – Mean linear slopes in basal area-based functional composition in Amazonia. As in Table 1 but showing the results for basal area, see Figure 1 for details.

Community	Potential size (cm y ⁻¹)	Water Deficit affiliation (mm y ⁻¹)	Wood Density (g cm ⁻³ y ⁻¹)
All community	0.03 (0.02 0.05)	0.01 (-0.04 0.07)	3x10 ⁻⁵ (-6x10 ⁻⁵ 1x10 ⁻⁴)
Gains (basal area)	0.06 (0.01 0.12)	-0.09 (-0.53 0.2)	-5x10 ⁻⁴ (-1x10 ⁻³ 1x10 ⁻⁴)
Gains (recruits)	0.06 (-0.09 0.19)	-0.08 (-0.7 0.6)	-2x10 ⁻⁴ (-2 x10 ⁻³ 2x10 ⁻³)
Losses	0.13 (-0.08 0.33)	-0.33 (-1.2 0.5)	-1x10 ⁻³ (-2x10 ⁻³ 3x10 ⁻⁴)
Net fluxes	-0.05 (-0.27 0.2)	0.24 (-0.7 1.19)	9x10 ⁻⁴ (-4x10 ⁻⁴ 2x10 ⁻³)

The influence of climate on functional composition

The trends in the intensity of extreme dry events ($MCWD_i$) within each plot were significantly correlated with trends in the losses of community basal-area WDA (Figure 4) and the fluxes of basal area WDA (Appendix S9). This indicates that the mortality of large wet-affiliated trees has increased in plots where the dry seasons have become more extreme. Trends in climate were also negatively correlated to losses of stem-based WD (Appendix S9). There was no significant correlation between trends in climate and trends in community weighted mean of any other trait or segments of the tree community (Appendix S9).




Figure 4 – Relationship between trends in climate and functional composition. The y-axis represents plot trends in water deficit affiliation (WDA) per year calculated as the linear slopes of basal area-based community weighted mean within the losses and x-axis shows the trends in most extreme dry season within a census interval ($MCWD_i$). The black line represents OLS linear regression and in the 1:1 line is shown in grey. Note that changes in the tree community are correlated to changes in climate (Kendall $\tau = -0.2$; $P < 0.01$), i.e., stronger drying trends favour the mortality of wet-affiliated taxa. Correlations hold when outliers are removed (Kendall $\tau = -0.4$; $P < 0.05$ when excluding outliers where climate trend > 5 mm y^{-1} and trends in losses > 10 mm y^{-1}).

Floristic trends

The floristic changes represented by the linear trends in abundance for individual genera are generally consistent with the functional changes observed. There has been a notable increase in the relative abundance of the dry-affiliated genera *Bertholletia*, *Brosimum* and *Pseudolmedia*, together with sharp decreases for the wet-affiliated *Aparisthmium*, *Fusaea*, *Inga* and *Mezilaurus*. Some large-statured genera have increased significantly, such as *Mora*, *Couratari*, and *Eschweilera*. A decrease in smaller-statured taxa, such as *Ouratea*, *Aniba*, *Marlierea* and *Astrocaryum* is also observed (Appendix S12). Palms (Arecaceae) have significantly declined in abundance (Appendix S12) overall, with marked declines of *Oenocarpus* and *Astrocaryum*, even though the hyperdominant palm genus *Euterpe* has significantly increased in abundance across the basin.

By analysing abundance trends of different taxa, it is possible to identify which genera contribute the most to the observed overall changes in functional composition (i.e., PS in basal area terms for the whole community and WDA stem density for recruits). The correlation between taxa PS and their trends in basal area was significant (Kendall $\tau = 0.2$; P-value < 0.01) and driven by an increase in some emergent and canopy genera such as *Eschweilera*, *Licania* and *Pouteria*, and a decrease in some sub-canopy and understorey genera such as *Iriarteia*, *Rinorea* and *Oenocarpus* (Figure 5 – Appendix S12). The decrease in WDA within the recruits was also explained by changes in floristic composition across the Amazon (Kendall $\tau = -0.16$; $P < 0.05$), with declines in the recruitment of new stems of wet affiliated Amazon genera such as *Mabea*, *Sterculia*, *Swartzia*, *Iryanthera* and *Theobroma* and recruitment increases for dry affiliates such as *Trema*, *Simarouba* and *Hieronyma* (Appendix S12).

Our analyses of the trends in abundance for the functional types defined for Amazonian trees by Fyllas *et al.* (2012) show small-statured non-pioneer taxa to be significantly decreasing

in abundance by 0.29% y^{-1} (Appendix S10). No significant trend was found for the other functional types (small-statured pioneers, tall pioneers, and tall non-pioneers). Note that this analysis was limited to the species classified by Fyllas *et al.* (2012) within these four functional types, which represent ca. 28% of the total number of individuals within our plots.




Figure 5 – Floristic changes behind the detected functional shifts. The y-axes represent mean linear slopes of plot level genera relative abundance across the Amazon in terms of number of individuals or basal area as a function of time, with each plots' contribution weighted by the square root of plot area and monitoring period. Grey horizontal lines indicate zero change. The x-axes represent genus-level traits. **(a)** Trends in relative basal area within the whole community vs. potential size (cm), plotted in the log scale to facilitate visualization; **(b)** trends in stem relative abundance

within the recruits vs. water deficit affiliation (mm). Genera that show confidence intervals not overlapping zero are shown in blue. Black contour marks the 10 most abundant genera in terms of number of stems.

Discussion

Here we report the first Amazon-wide analyses of temporal trends in functional and floristic composition of lowland tree communities across 106 Amazonian inventory plots analysed over three decades. We find evidence of climate-induced shifts in community composition. Recruits have become more dry-affiliated (Table 1) and the mortality of wet-affiliated trees has increased in the areas where the drying trend was stronger (Figure 4), suggesting a direct effect of climate change on functional composition. This compositional shift is consistent with the detected intensification of the seasonal drought across the majority of permanent monitoring plots in Amazonia. The trends in potential size (PS) and wood density (WD) further indicate that changes within the Amazon forest are likely to be a consequence of a combination of drivers. However, our data also highlight the relative inertia of tropical forest communities in their ability to respond to environmental changes. For instance, the detected change in tree communities, with recruits becoming more dry-affiliated by 0.45 mm y^{-1} , is substantially smaller in magnitude when compared to the climate trend of 1.1 mm y^{-1} .

The functional shifts observed here are mirrored by underlying floristic changes when our data are analysed in terms of discrete taxonomic units (Figure 5). The genus-level floristic shifts behind the significant changes detected (Table 1 and 2) illustrates how functional responses result from actual floristic change across Amazonian communities. However, the relationships between traits and floristic shifts also show significant scatter, suggesting idiosyncratic responses of each taxon and the complexity of this highly diverse system.

Climate-induced changes in floristic composition

We detected an increase in the abundance of drought-tolerant genera across Amazonia when analysing the recruitment and mortality within tree communities (Table 1, Figure 4), consistent with the hypothesis that tree communities are responding to the changes in climate. Our large-scale results are consistent with tree community shifts towards more drought-tolerant taxa reported in some neotropical (Butt *et al.*, 2014, Enquist & Enquist, 2011, Feeley *et al.*, 2011b) and west African forest localities (Fauset *et al.*, 2012), and some temperate localities (McIntyre *et al.*, 2015). Across Amazonia we find that greater mortality of wet-affiliated taxa over time is related to the degree to which water-stress has increased within each community, providing direct evidence of the influence of climate on community dynamics (Figure 4). This only emerged when analysing trends in basal area, indicating that it is driven by the increase in mortality of large wet-affiliated trees. Indeed, large trees are expected to suffer the most under drought conditions by being more vulnerable to embolism, and thus more likely to die from hydraulic failure under drought-stress (McDowell & Allen, 2015, Rowland *et al.*, 2015). In addition, further supporting the conclusion that this reflects concerted, widespread changes in Amazon forest mortality, we also found an increase in the potential size of the dead trees when analyses were repeated considering each cluster of neighbouring plots as a single sample unit (Appendix S7). Somewhat surprisingly we find losses in stem-based WD to be negatively correlated to changes in climate (Appendix S9), which suggests that WD may not be a good proxy of drought vulnerability.

Nonetheless our results suggest that other drivers are also causing compositional shifts in Amazon forests. For example, although we did observe an increase in mortality of large, wet affiliated trees, consistent with the effect of the 2005 drought (Phillips *et al.*, 2009), somewhat surprisingly our analyses also show a slight increase in the basal area of potentially large tree

genera across the basin. These results contradict the expectations that droughts would preferably impact larger trees but are in line with the suggestion that smaller-statured trees are more vulnerable to droughts as they have shallower roots than larger-statured trees (Condit *et al.*, 1996, Fauset *et al.*, 2012, Wright, 1992). Most likely, it appears that the increase in mortality of large trees in some areas of the basin (Figure 4, Appendix S9) is a consequence of drought events such as the 2005 drought (cf. Bennett *et al.*, 2015, Phillips *et al.*, 2010), but that it this mortality has been insufficient to counter a longer-term tendency towards increased basal area of large-statured taxa across Amazonia.

Additional drivers of compositional change

Larger trees have greater competitive capacity and are anticipated to gain disproportionately with additional resources (Coomes *et al.*, 2011, Enquist *et al.*, 2009, Enquist *et al.*, 1999). The increase of atmospheric CO₂ potentially provides a parsimonious explanation for the observed relative increase in large-tree genera (Table 2). If the increase in atmospheric CO₂ of ca. 5% per decade since the 1980s (Tans, 2016) is stimulating plant growth or increasing water-use efficiency, then taxa that tend to compete better for light, notably larger trees (Poorter *et al.*, 2005), are likely to gain a greater competitive advantage (Falster & Westoby, 2003). Although further investigation is needed to confirm this hypothesis, our results show an increase in basal area of large-statured taxa (Figure 3; Table 2) and a decrease in abundance of non-pioneer small-statured taxa (Appendix S10), both consistent with the expectations from increased competition. Additionally, the relative increase of larger genera is also observed in the broader dataset including the fringes of Amazonia (*Extended Amazonia* see Supplementary material), which supports the inference of a ubiquitous driver of compositional change. Our observations are also in line with a series of stand-level studies that show increases in above ground biomass, growth, mortality and recruitment across Amazonia – all trends expected as

an outcome from increased atmospheric CO₂ (Brienen *et al.*, 2015, Phillips & Gentry, 1994, Phillips *et al.*, 1998).

While considerable interest has focussed on the atmospheric and climatic changes, particularly drought, other environmental changes may be important. Conceivably one or more of these could have pervasive impacts on mature forests across the basin to rival or exceed any climate impacts. Amazonian forests have been hypothesized to be recovering from previous disturbances that are either natural (Chambers *et al.*, 2013, van der Sande *et al.*, 2016) or human-driven, particularly in pre-Colombian times (McMichael *et al.*, 2017). If this is the case, we would expect the forest to be following a successional trajectory characterised by a shift from pioneers (low wood density) to mature forest species (high wood density) (e.g., Chave *et al.*, 2008, Connell & Slatyer, 1977, Lewis *et al.*, 2009a). While the observed relative increase in basal area of larger taxa is consistent with widespread recovery from disturbance (Chave *et al.*, 2008, Wright, 2005), the 106 Amazonian inventory plots show no significant shift in wood density (WD) across the whole community, or perhaps more importantly, among the recruits (Table 2). We note that our WD data (Chave *et al.*, 2009, Zanne *et al.*, 2009) provide a less than perfect insight into actual change given that there is likely to be additional spatial and intra-generic variation (Baraloto *et al.*, 2010, Patiño *et al.*, 2009) that we cannot account for. However, WD is considerably conserved across the phylogeny, and genus-level wood density has been found to be adequate to distinguish between late successional and pioneer genera (Coelho de Souza *et al.*, 2016). Moreover the only functional group that has clearly *lost* ground over our monitoring window are the short-statured non-pioneers which are the best adapted to low light, late-successional environments (Appendix S10). Compositional analyses further suggest that some pioneers are increasing, most notably an increase of 3.4% ha⁻¹y⁻¹ in the relative abundance of *Cecropia*, a key early successional taxon (Appendix S12), which

typically dominates in early stages of succession but declines at later stages of the successional trajectory (Zalamea *et al.*, 2012). The increase in the abundance of light-demanding taxa may be a consequence of an acceleration in the canopy gap dynamics caused by the increase in baseline mortality rates (Baker *et al.*, 2016, Brien *et al.*, 2015). Finally, if these forest plots are recovering from the impact of Amazonian peoples who favoured especially useful species, then we might expect domesticated taxa *sensu* Levis *et al.* (2017) to now be decreasing in abundance following the relaxation of this anthropogenic influence. No such trend is observed in our data (Appendix S11).

There has been considerable concern regarding the ecosystem impacts of widespread removal of large-bodied frugivores. In particular, it has been repeatedly suggested that hunting will or may have already altered tree composition in tropical forests (e.g. Doughty *et al.*, 2016, Osuri *et al.*, 2016, Peres *et al.*, 2016, Peres & Palacios, 2007, Terborgh *et al.*, 2008) via dispersal failure of zoochoric trees (Chapman & Chapman, 1995). These tend to have high wood density and large stature, so a recruitment failure is predicted to lead to a reduction in the prevalence of both of these traits and thus in Amazonian biomass (Bello *et al.*, 2015, Peres *et al.*, 2016). This study was designed to understand floristic dynamics in intact Amazonian forests and not to evaluate the effects of hunting pressure, which is likely to more strongly affect areas adjacent to rural communities, roads and rivers (Peres & Lake, 2003). However, the increase in potential size and the lack of change in wood density within the recruits (Figure 3) contradict expectations of the hunting hypothesis as a driver of recent community composition change in intact forests, which, again, does not rule out the possibility of hunting pressure causing shifts in composition in particular locations where this activity is stronger.

The pace of change

Changes in tree communities are expected to substantially lag behind environmental changes as trees are sessile and long-lived (Blonder *et al.*, 2017, Davis, 1989, Hubbell, 2001, Lenoir *et al.*, 2008, Svenning & Sandel, 2013). Our results are consistent with this prediction. In other systems where climate gradients are almost unidirectional it is possible to assess the speed at which communities are expected to be moving across spaces to follow climate (Devictor *et al.*, 2012), but this is not the case for precipitation in Amazonia where precipitation regime is heterogeneous at multiple spatial scales. However, by ensuring that climate and water-deficit affiliation are calculated in the same scale we can compare the degree to which climate and communities are changing. Across Amazonia, plots have undergone an average drying trend of -1.1 mm y^{-1} MCWD (Figure 2). Notably, change in tree communities did not keep pace with the change in climate - the mean trend in water-deficit affiliation for the whole community is two orders of magnitude smaller (0.01 mm y^{-1} , Table 1), with confidence intervals overlapping zero. However, a much shorter lag is observed when analysing the net fluxes, i.e., recruits – deaths (-0.45 mm y^{-1} , Table 1), indicating that although responses are slow, this system has some dynamic capacity to respond to changes in climate.

The observed pace of change is a reflection of the nature of these communities. Recruitment and mortality of trees $\geq 10 \text{ cm D}$ averaged 9 trees per hectare per year between 1985 and 2010 across the basin (Brienen *et al.*, 2015). Considering that in our data mean stem density is $520 \text{ trees ha}^{-1}$ and mean plot-monitoring length is 14 years, we can expect by the final census an accumulated turnover of $\approx 24\%$ of stems. Thus, we should expect *a priori* that whole community-level composition would be only weakly affected by changes over the measurement period and would instead largely reflecting legacy effects of recruitment and mortality processes occurring decades prior to the onset of the monitoring period (Davis, 1989).

Our results provide empirical evidence of the inertia within this system and clearly raise concerns about whether forests here will be able to track further climate change anticipated over coming decades.

This study provides the first Amazon-wide picture of functional and floristic dynamics over the last 30 years. Models have predicted a strengthening of the dry season over the Amazon (Boisier *et al.*, 2015, Joetzjer *et al.*, 2013), and an increase in water-stress as a consequence of rising temperature (Pokhrel *et al.*, 2014). But there have been few attempts to account for changes in composition, which may modulate the whole forest ecological impact of ecophysiological drivers such as increasing vapour pressure deficit (Levine *et al.*, 2016, Sakschewski *et al.*, 2016). The velocity and the magnitude of compositional changes presented here should be considered in attempts to model the dynamics of these forests under future climate. Our results show that a slow shift towards a more dry-affiliated Amazonia is underway. If such a floristic shift is substantial enough to increase the resilience of Amazon tree communities to future droughts, it will still come with a price in terms of tree biodiversity, since wet-affiliated restricted taxa represent the majority of Amazonian tree flora (Esquivel-Muelbert *et al.*, 2017a). Furthermore, although our results demonstrate that changes in composition are possible, the inertia intrinsic to these communities means that they will substantially lag behind climate change. Droughts are continuing to impact the basin (Erfanian *et al.*, 2017, Jiménez-Muñoz *et al.*, 2016). If this lag persists, intact Amazonian forests may be increasingly dominated by sub-optimally adapted trees in terms of their preferred climate space, potentially threatening the ability of these ecosystems to provide key services such as protecting biodiversity and sequestering and storing carbon.

Acknowledgements

This paper is a product of the RAINFOR network. It integrates the effort of hundreds of researchers and field assistants across Amazonia over the last three decades and the sustained support from rural communities and local institutions. We thank the following individuals in particular Atila Alves de Oliveira, Benoit Burban, Bert van Ulf, Christopher Baraloto, Eliana Riascos, Foster Brown, Francisco Gómez, Freddy Ramirez Arevalo, Iêda Leão do Amaral, Irina Mendoza Polo, Jean Olivier, Joey Talbot, John Lleigue, Marcela Serna, Michel Baisie, Lilian Blanc, Patricia Alvarez Loayza, Roderick Zagt, Nadir Pallqui Camacho, Nazareno Martinez, Olaf Banki, Pétrus Naisso, Samaria Murakami, Samuel Almeida, Timothy Killeen, Tomas Dario Gutierrez, Walter Huaraca Huasco, Wemo Betian and Vincent Bezard. Support for RAINFOR has come from the Natural Environment Research Council (NERC) Urgency Grants and NERC Consortium Grants “AMAZONICA” (NE/F005806/1), “TROBIT” (NE/D005590/1) and “BIO-RED” (NE/N012542/1), a European Research Council (ERC) grant (T-FORCES, “Tropical Forests in the Changing Earth System”), the Gordon and Betty Moore Foundation, the European Union’s Seventh Framework Programme (282664, “AMAZALERT”) and the Royal Society (CH160091). O.L.P. was supported by an ERC Advanced Grant and a Royal Society Wolfson Research Merit Award. K.G.D. was supported by a Leverhulme Trust International Academic Fellowship. This paper is part of the PhD of A.E-M, which was funded by the ERC T-FORCES grant. A.E-M. is currently supported by T-FORCES and the NERC project “TREMOR” (NE/N004655/1).

References

- Allen RG, Smith M, Pereira LS, Perrier A (1994) An update for the calculation of reference evapotranspiration. *ICID Bulletin*, **43**, 35-92.
- Aragao L, Malhi Y, Roman-Cuesta RM, Saatchi S, Anderson LO, Shimabukuro YE (2007) Spatial patterns and fire response of recent Amazonian droughts. *Geophysical Research Letters*, **34**.
- Baker TR, Phillips OL, Malhi Y *et al.* (2004) Variation in wood density determines spatial patterns in Amazonian forest biomass. *Global Change Biology*, **10**, 545-562.
- Baker TR, Vela Díaz DM, Chama Moscoso V *et al.* (2016) Consistent, small effects of treefall disturbances on the composition and diversity of four Amazonian forests. *Journal of Ecology*, **104**, 497-506.
- Baraloto C, Paine CET, Patino S, Bonal D, Herault B, Chave J (2010) Functional trait variation and sampling strategies in species-rich plant communities. *Functional Ecology*, **24**, 208-216.
- Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. 2015, **67**, 48.
- Bello C, Galetti M, Pizo MA *et al.* (2015) Defaunation affects carbon storage in tropical forests. *Science Advances*, **1**.
- Bennett AC, Mcdowell NG, Allen CD, Anderson-Teixeira KJ (2015) Larger trees suffer most during drought in forests worldwide. *Nature Plants*, **1**, 15139.
- Blonder B, Moulton DE, Blois J *et al.* (2017) Predictability in community dynamics. *Ecology Letters*, **20**, 293-306.
- Boisier JP, Ciais P, Ducharne A, Guimberteau M (2015) Projected strengthening of Amazonian dry season by constrained climate model simulations. *Nature Climate Change*, **5**, 656-660.
- Bowler DE, Hof C, Haase P *et al.* (2017) Cross-realm assessment of climate change impacts on species' abundance trends. *Nat Ecol Evol*, **1**, 67.
- Brienen RJW, Phillips OL, Feldpausch TR *et al.* (2015) Long-term decline of the Amazon carbon sink. *Nature*, **519**, 344-348.
- Butt N, Malhi Y, New M *et al.* (2014) Shifting dynamics of climate-functional groups in old-growth Amazonian forests. *Plant Ecology & Diversity*, **7**, 267-279.
- Cardoso D, Särkinen T, Alexander S *et al.* (2017) Amazon plant diversity revealed by a taxonomically verified species list. *Proceedings of the National Academy of Sciences*, **114**, 10695-10700.
- Chambers JQ, Negron-Juarez RI, Marra DM *et al.* (2013) The steady-state mosaic of disturbance and succession across an old-growth Central Amazon forest landscape. *Proceedings of the National Academy of Sciences*, **110**, 3949-3954.
- Chapman CA, Chapman LJ (1995) Survival without dispersers: seedling recruitment under parents. *Conservation Biology*, **9**, 675-678.

711 Chave J, Condit R, Muller-Landau HC *et al.* (2008) Assessing evidence for a pervasive alteration in
712 tropical tree communities. *Plos Biology*, **6**, 455-462.

713 Chave J, Coomes D, Jansen S, Lewis SL, Swenson NG, Zanne AE (2009) Towards a worldwide wood
714 economics spectrum. *Ecology Letters*, **12**, 351-366.

715 Chen IC, Shiu HJ, Benedick S *et al.* (2009) Elevation increases in moth assemblages over 42 years on a
716 tropical mountain. *Proceedings of the National Academy of Sciences of the United States of*
717 *America*, **106**, 1479-1483.

718 Coelho De Souza F, Dexter KG, Phillips OL *et al.* (2016) Evolutionary heritage influences Amazon tree
719 ecology. *Proceedings of the Royal Society B: Biological Sciences*, **283**.

720 Condit R, Hubbell SP, Foster RB (1996) Changes in tree species abundance in a Neotropical forest:
721 Impact of climate change. *Journal of Tropical Ecology*, **12**, 231-256.

722 Connell JH, Slatyer RO (1977) Mechanisms of succession in natural communities and their role in
723 community stability and organization. *American Naturalist*, **111**, 1119-1144.

724 Conway TJ, Tans PP (2016) Trends in atmospheric carbon dioxide. pp Page.

725 Coomes DA, Lines ER, Allen RB (2011) Moving on from Metabolic Scaling Theory: hierarchical models
726 of tree growth and asymmetric competition for light. *Journal of Ecology*, **99**, 748-756.

727 Davis MB (1989) Lags in vegetation response to greenhouse warming. *Climatic Change*, **15**, 75-82.

728 Devictor V, Van Swaay C, Brereton T *et al.* (2012) Differences in the climatic debts of birds and
729 butterflies at a continental scale. *Nature Climate Change*, **2**, 121.

730 Diaz S, Lavorel S, De Bello F, Quetier F, Grigulis K, Robson M (2007) Incorporating plant functional
731 diversity effects in ecosystem service assessments. *Proceedings of the National Academy of*
732 *Sciences of the United States of America*, **104**, 20684-20689.

733 Doughty CE, Wolf A, Morueta-Holme N *et al.* (2016) Megafauna extinction, tree species range
734 reduction, and carbon storage in Amazonian forests. *Ecography*, **39**, 194-203.

735 Duffy PB, Brando P, Asner GP, Field CB (2015) Projections of future meteorological drought and wet
736 periods in the Amazon. *Proceedings of the National Academy of Sciences of the United States*
737 *of America*, **112**.

738 Duque A, Stevenson PR, Feeley KJ (2015) Thermophilization of adult and juvenile tree communities in
739 the northern tropical Andes. *Proceedings of the National Academy of Sciences*, **112**, 10744-
740 10749.

741 Engelbrecht BMJ, Comita LS, Condit R, Kursar TA, Tyree MT, Turner BL, Hubbell SP (2007) Drought
742 sensitivity shapes species distribution patterns in tropical forests. *Nature*, **447**, 80-U82.

743 Enquist BJ, Enquist CaF (2011) Long-term change within a Neotropical forest: assessing differential
744 functional and floristic responses to disturbance and drought. *Global Change Biology*, **17**,
745 1408-1424.

746 Enquist BJ, West GB, Brown JH (2009) Extensions and evaluations of a general quantitative theory of
747 forest structure and dynamics. *Proceedings of the National Academy of Sciences*, **106**, 7046-
748 7051.

749 Enquist BJ, West GB, Charnov EL, Brown JH (1999) Allometric scaling of production and life-history
750 variation in vascular plants. *Nature*, **401**, 907-911.

751 Erfanian A, Wang G, Fomenko L (2017) Unprecedented drought over tropical South America in 2016:
752 significantly under-predicted by tropical SST. *Scientific Reports*, **7**, 5811.

753 Espinoza JC, Marengo JA, Ronchail J, Carpio JM, Flores LN, Guyot JL (2014) The extreme 2014 flood in
754 south-western Amazon basin: the role of tropical-subtropical South Atlantic SST gradient.
755 *Environmental Research Letters*, **9**, 124007.

756 Esquivel-Muelbert A, Baker TR, Dexter KG *et al.* (2017a) Seasonal drought limits tree species across
757 the Neotropics. *Ecography*, **40**, 618-629.

758 Esquivel-Muelbert A, Galbraith D, Dexter KG *et al.* (2017b) Biogeographic distributions of neotropical
759 trees reflect their directly measured drought tolerances. *Scientific Reports*, **7**, 8334.

760 Falster DS, Westoby M (2003) Plant height and evolutionary games. *Trends in Ecology & Evolution*, **18**,
761 337-343.

762 Falster DS, Westoby M (2005) Alternative height strategies among 45 dicot rain forest species from
763 tropical Queensland, Australia. *Journal of Ecology*, **93**, 521-535.

764 Fauset S, Baker TR, Lewis SL *et al.* (2012) Drought-induced shifts in the floristic and functional
765 composition of tropical forests in Ghana. *Ecology Letters*, **15**, 1120-1129.

766 Fauset S, Johnson MO, Gloor M *et al.* (2015) Hyperdominance in Amazonian forest carbon cycling.
767 *Nature Communications*, **6**.

768 Feeley KJ, Davies SJ, Perez R, Hubbell SP, Foster RB (2011a) Directional changes in the species
769 composition of a tropical forest. *Ecology*, **92**, 871-882.

770 Feeley KJ, Silman MR, Bush MB *et al.* (2011b) Upslope migration of Andean trees. *Journal of*
771 *Biogeography*, **38**, 783-791.

772 Feldpausch TR, Lloyd J, Lewis SL *et al.* (2012) Tree height integrated into pantropical forest biomass
773 estimates. *Biogeosciences*, **9**, 3381-3403.

774 Feldpausch TR, Phillips OL, Brien RJW *et al.* (2016) Amazon forest response to repeated droughts.
775 *Global Biogeochemical Cycles*, **30**, 964-982.

776 Fisher JI, Hurtt GC, Thomas RQ, Chambers JQ (2008) Clustered disturbances lead to bias in large-scale
777 estimates based on forest sample plots. *Ecology Letters*, **11**, 554-563.

778 Flores O, Coomes DA (2011) Estimating the wood density of species for carbon stock assessments.
779 *Methods in Ecology and Evolution*, **2**, 214-220.

780 Francis AP, Currie DJ (2003) A globally consistent richness-climate relationship for angiosperms.
781 *American Naturalist*, **161**, 523-536.

782 Fyllas NM, Patino S, Baker TR *et al.* (2009) Basin-wide variations in foliar properties of Amazonian
783 forest: phylogeny, soils and climate. *Biogeosciences*, **6**, 2677-2708.

784 Fyllas NM, Quesada CA, Lloyd J (2012) Deriving Plant Functional Types for Amazonian forests for use
785 in vegetation dynamics models. *Perspectives in Plant Ecology, Evolution and Systematics*, **14**,
786 97-110.

787 Gatti LV, Gloor M, Miller JB *et al.* (2014) Drought sensitivity of Amazonian carbon balance revealed by
788 atmospheric measurements. *Nature*, **506**, 76-80.

789 Gentry AH (1988) Changes in plant community diversity and floristic composition on environmental
790 and geographical gradients. *Annals of the Missouri Botanical Garden*, **75**, 1-34.

791 Giardina F, Konings AG, Kennedy D, Alemohammad SH, Oliveira RS, Uriarte M, Gentine P (2018) Tall
792 Amazonian forests are less sensitive to precipitation variability. *Nature Geoscience*, **11**, 405-
793 409.

794 Gloor M, Barichivich J, Ziv G *et al.* (2015) Recent Amazon Climate as background for possible ongoing
795 and future changes of Amazon humid forests. *Global Biogeochemical Cycles*, **29**, 1384-1399.

796 Harris I, Jones PD, Osborn TJ, Lister DH (2014) Updated high-resolution grids of monthly climatic
797 observations – the CRU TS3.10 Dataset. *International Journal of Climatology*, **34**, 623-642.

798 Hilker T, Lyapustin AI, Tucker CJ *et al.* (2014) Vegetation dynamics and rainfall sensitivity of the
799 Amazon. *Proceedings of the National Academy of Sciences*, **111**, 16041-16046.

800 Hubbell SP (2001) *The Unified Neutral Theory of Biodiversity and Biogeography*, Princeton,
801 Princeton University Press.

802 Huffman GJ, Bolvin DT, Nelkin EJ *et al.* (2007) The TRMM Multisatellite Precipitation Analysis (TMPA):
803 Quasi-Global, Multiyear, Combined-Sensor Precipitation Estimates at Fine Scales. *Journal of*
804 *Hydrometeorology*, **8**, 38-55.

805 Jiménez-Muñoz JC, Mattar C, Barichivich J *et al.* (2016) Record-breaking warming and extreme drought
806 in the Amazon rainforest during the course of El Niño 2015–2016. *Scientific Reports*, **6**, 33130.

807 Jiménez-Muñoz JC, Sobrino JA, Mattar C, Malhi Y (2013) Spatial and temporal patterns of the recent
808 warming of the Amazon forest. *Journal of Geophysical Research: Atmospheres*, **118**, 5204-
809 5215.

810 Joetzjer E, Douville H, Delire C, Ciais P (2013) Present-day and future Amazonian precipitation in global
811 climate models: CMIP5 versus CMIP3. *Climate Dynamics*, **41**, 2921-2936.

812 Laurance WF (2004) Forest-climate interactions in fragmented tropical landscapes. *Philosophical*
813 *Transactions of the Royal Society of London Series B-Biological Sciences*, **359**, 345-352.

814 Laurance WF, Oliveira AA, Laurance SG *et al.* (2004) Pervasive alteration of tree communities in
815 undisturbed Amazonian forests. *Nature*, **428**, 171-175.

816 Lenoir J, Gegout JC, Marquet PA, De Ruffray P, Brisse H (2008) A significant upward shift in plant
817 species optimum elevation during the 20th century. *Science*, **320**, 1768-1771.

818 Levine NM, Zhang K, Longo M *et al.* (2016) Ecosystem heterogeneity determines the ecological
819 resilience of the Amazon to climate change. *Proceedings of the National Academy of Sciences*,
820 **113**, 793-797.

821 Levis C, Costa FRC, Bongers F *et al.* (2017) Persistent effects of pre-Columbian plant domestication on
822 Amazonian forest composition. *Science*, **355**, 925-931.

823 Lewis SL, Brando PM, Phillips OL, Van Der Heijden GMF, Nepstad D (2011) The 2010 Amazon drought.
824 *Science*, **331**, 554-554.

825 Lewis SL, Lloyd J, Sitch S, Mitchard ETA, Laurance WF (2009a) Changing ecology of tropical forests:
826 evidence and drivers. In: *Annual Review of Ecology Evolution and Systematics*. pp Page. Palo
827 Alto, Annual Reviews.

828 Lewis SL, Lopez-Gonzalez G, Sonke B *et al.* (2009b) Increasing carbon storage in intact African tropical
829 forests. *Nature*, **457**, 1003-U1003.

830 Lloyd J, Farquhar GD (1996) The CO₂ dependence of photosynthesis, plant growth responses to
831 elevated atmospheric CO₂ concentrations and their interaction with soil nutrient status .1.
832 General principles and forest ecosystems. *Functional Ecology*, **10**, 4-32.

833 Lloyd J, Farquhar GD (2008) Effects of rising temperatures and CO₂ on the physiology of tropical forest
834 trees. *Philosophical Transactions of the Royal Society B-Biological Sciences*, **363**, 1811-1817.

835 Lopez-Gonzalez G, Lewis SL, Burkitt M, Phillips OL (2011) ForestPlots.net: a web application and
836 research tool to manage and analyse tropical forest plot data. *Journal of Vegetation Science*,
837 **22**, 610-613.

838 Malhi Y, Phillips OL, Lloyd J *et al.* (2002) An international network to monitor the structure,
839 composition and dynamics of Amazonian forests (RAINFOR). *Journal of Vegetation Science*,
840 **13**, 439-450.

841 Marengo JA, Espinoza JC (2016) Extreme seasonal droughts and floods in Amazonia: causes, trends
842 and impacts. *International Journal of Climatology*, **36**, 1033-1050.

843 Marengo JA, Nobre CA, Tomasella J *et al.* (2008) The drought of Amazonia in 2005. *Journal of Climate*,
844 **21**, 495-516.

845 Marengo JA, Tomasella J, Alves LM, Soares WR, Rodriguez DA (2011) The drought of 2010 in the
846 context of historical droughts in the Amazon region. *Geophysical Research Letters*, **38**.

847 Mcdowell NG, Allen CD (2015) Darcy's law predicts widespread forest mortality under climate
848 warming. *Nature Clim. Change*, **5**, 669-672.

849 McGill BJ, Enquist BJ, Weiher E, Westoby M (2006) Rebuilding community ecology from functional
850 traits. *Trends in Ecology & Evolution*, **21**, 178-185.

851 McIntyre PJ, Thorne JH, Dolanc CR, Flint AL, Flint LE, Kelly M, Ackerly DD (2015) Twentieth-century
852 shifts in forest structure in California: Denser forests, smaller trees, and increased dominance
853 of oaks. *Proceedings of the National Academy of Sciences*, **112**, 1458-1463.

854 Mcmichael CNH, Matthews-Bird F, Farfan-Rios W, Feeley KJ (2017) Ancient human disturbances may
855 be skewing our understanding of Amazonian forests. *Proceedings of the National Academy of*
856 *Sciences*, **114**, 522-527.

857 Muller-Landau HC (2004) Interspecific and inter-site variation in wood specific gravity of tropical trees.
858 *Biotropica*, **36**, 20-32.

859 Nascimento HEM, Laurance WF, Condit R, Laurance SG, D'angelo S, Andrade AC (2005) Demographic
860 and life-history correlates for Amazonian trees. *Journal of Vegetation Science*, **16**, 625-634.

861 Nepstad DC, Tohver IM, Ray D, Moutinho P, Cardinot G (2007) Mortality of large trees and lianas
862 following experimental drought in an amazon forest. *Ecology*, **88**, 2259-2269.

863 Osuri AM, Ratnam J, Varma V *et al.* (2016) Contrasting effects of defaunation on aboveground carbon
864 storage across the global tropics. *Nat Commun*, **7**.

865 Pan YD, Birdsey RA, Fang JY *et al.* (2011) A Large and Persistent Carbon Sink in the World's Forests.
866 *Science*, **333**, 988-993.

867 Patiño S, Lloyd J, Paiva R *et al.* (2009) Branch xylem density variations across the Amazon Basin.
868 *Biogeosciences*, **6**, 545-568.

869 Penone C, Davidson AD, Shoemaker KT *et al.* (2014) Imputation of missing data in life-history trait
870 datasets: which approach performs the best? *Methods in Ecology and Evolution*, **5**, 961-970.

871 Peres CA, Emilio T, Schietti J, Desmoulière SJM, Levi T (2016) Dispersal limitation induces long-term
872 biomass collapse in overhunted Amazonian forests. *Proceedings of the National Academy of*
873 *Sciences*, **113**, 892-897.

874 Peres CA, Lake IR (2003) Extent of nontimber resource extraction in tropical forests: Accessibility to
875 game vertebrates by hunters in the Amazon basin. *Conservation Biology*, **17**, 521-535.

876 Peres CA, Palacios E (2007) Basin-wide effects of game harvest on vertebrate population densities in
877 Amazonian forests: implications for animal-mediated seed dispersal. *Biotropica*, **39**, 304-315.

878 Phillips O, Baker T, Feldpausch T *et al.* (2016) RAINFOR field manual for plot establishment and
879 remeasurement.

880 Phillips OL, Aragao L, Lewis SL *et al.* (2009) Drought sensitivity of the amazon rainforest. *Science*, **323**,
881 1344-1347.

882 Phillips OL, Gentry AH (1994) Increasing turnover through time in tropical forests. *Science*, **263**, 954-
883 958.

884 Phillips OL, Malhi Y, Higuchi N *et al.* (1998) Changes in the Carbon Balance of Tropical Forests: Evidence
885 from Long-Term Plots. *Science*, **282**, 439-442.

886 Phillips OL, Van Der Heijden G, Lewis SL *et al.* (2010) Drought-mortality relationships for tropical
887 forests. *New Phytologist*, **187**, 631-646.

888 Pitman NCA, Terborgh JW, Silman MR *et al.* (2002) A comparison of tree species diversity in two upper
889 Amazonian forests. *Ecology*, **83**, 3210-3224.

890 Pokhrel YN, Fan Y, Miguez-Macho G (2014) Potential hydrologic changes in the Amazon by the end of
891 the 21st century and the groundwater buffer. *Environmental Research Letters*, **9**.

892 Poorter L, Bongers F, Sterck FJ, Wöll H (2005) Beyond the regeneration phase: differentiation of
893 height–light trajectories among tropical tree species. *Journal of Ecology*, **93**, 256-267.

894 Poorter L, Mcdonald I, Alarcon A *et al.* (2010) The importance of wood traits and hydraulic
895 conductance for the performance and life history strategies of 42 rainforest tree species. *New*
896 *Phytologist*, **185**, 481-492.

897 Qie L, Lewis SL, Sullivan MJP *et al.* (2017) Long-term carbon sink in Borneo’s forests halted by drought
898 and vulnerable to edge effects. *Nature Communications*.

899 Rowland L, Da Costa ACL, Galbraith DR *et al.* (2015) Death from drought in tropical forests is triggered
900 by hydraulics not carbon starvation. *Nature*, 459–464.

901 Sakschewski B, Von Bloh W, Boit A *et al.* (2016) Resilience of Amazon forests emerges from plant trait
902 diversity. *Nature Clim. Change*, **6**, 1032-1036.

903 Stahl C, Hérault B, Rossi V, Burban B, Bréchet C, Bonal D (2013) Depth of soil water uptake by tropical
904 rainforest trees during dry periods: does tree dimension matter? *Oecologia*, **173**, 1191-1201.

905 Svenning J-C, Sandel B (2013) Disequilibrium vegetation dynamics under future climate change.
906 *American Journal of Botany*.

907 Tans P (2016) Trends in atmospheric carbon dioxide. pp Page.

908 Ter Steege H, Nigel CA, Sabatier D *et al.* (2013) Hyperdominance in the amazonian tree flora. *Science*,
909 **342**, 325-+.

910 Terborgh J, Nunez-Iturri G, Pitman NCA *et al.* (2008) Tree recruitment in an empty forest. *Ecology*, **89**,
911 1757-1768.

912 Thomas CD, Cameron A, Green RE *et al.* (2004) Extinction risk from climate change. *Nature*, **427**, 145-
913 148.

914 Van Der Sande MT, Arets EJMM, Peña-Claros M *et al.* (2016) Old-growth Neotropical forests are
915 shifting in species and trait composition. *Ecological Monographs*, n/a-n/a.

916 Van Der Sleen P, Groenendijk P, Vlam M *et al.* (2015) No growth stimulation of tropical trees by 150
917 years of CO2 fertilization but water-use efficiency increased. *Nature Geosci*, **8**, 24-28.

918 Violle C, Reich PB, Pacala SW, Enquist BJ, Kattge J (2014) The emergence and promise of functional
919 biogeography. *Proceedings of the National Academy of Sciences of the United States of*
920 *America*, **111**, 13690-13696.

921 Wright SJ (1992) Seasonal drought, soil fertility and the species density of tropical forest plant
922 communities. *Trends in Ecology & Evolution*, **7**, 260-263.

923 Wright SJ (2005) Tropical forests in a changing environment. *Trends in Ecology & Evolution*, **20**, 553-
924 560.

925 Zalamea P-C, Heuret P, Sarmiento C *et al.* (2012) The Genus Cecropia: A Biological Clock to Estimate
926 the Age of Recently Disturbed Areas in the Neotropics. Plos One, **7**, e42643.

927 Zanne AE, Lopez-Gonzalez G, Coomes DA *et al.* (2009) Data from: Towards a worldwide wood
928 economics spectrum. pp Page, Dryad Data Repository.

929 Zhu Z, Piao S, Myneni RB *et al.* (2016) Greening of the Earth and its drivers. Nature Climate Change, **6**,
930 791.

931 Zuleta D, Duque A, Cardenas D, Muller-Landau HC, Davies S (2017) Drought-induced mortality patterns
932 and rapid biomass recovery in a terra firme forest in the Colombian Amazon. Ecology, **98**,
933 2538-2546.

934